



Applications of Gauss's Law

What lies behind us and what lies before us are
tiny matters compared to what lies within us.

Emerson

Don't walk before me, I may not follow; Don't
walk behind me, I may not lead; Walk beside
me and be my friend.


$$\overline{\Phi_E} = E \cdot A ; \quad \overline{E} = \oint E \cdot dA = \frac{q_{in}}{\epsilon_0}$$

Symmetric Charge Distribution

- We want to select a surface such that one or more of the following conditions are met:
 1. The value of the E-field can be argued by symmetry to be constant over the surface.
 2. The dot product can be expressed as a simple algebraic product $\overbrace{E} \cdot dA$, because **E** and **dA** are parallel. $E \parallel dA \Rightarrow E \cdot dA = E dA$
 3. The dot product is zero b/c **E** and **dA** are perpendicular. $E \perp dA \Rightarrow E \cdot dA = 0$
 4. The field can be argued to be zero over the surface.

Examples, pages 750 - 754


1. Calculated the electric field due to a point charge, $+q$. $E \perp dA$




$$\oint E \cdot dA = \frac{q_{in}}{\epsilon_0}$$
$$E (4\pi r^2) = \frac{q}{\epsilon_0}$$
$$E = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2}$$
$$E = \frac{kq}{r^2}$$

2. An insulating solid sphere of radius R has a uniform volume charge density ρ and carries a total positive charge Q .

a) Calculate E-field outside sphere.



b) Calculate E-field inside sphere.



a) $r > R$, $E = \frac{kQ}{r^2}$

b) $r < R$, $\oint E \cdot dA = \frac{q_{in}}{\epsilon_0}$

$$E (4\pi r^2) = \frac{\rho \frac{4}{3}\pi r^3}{\epsilon_0}$$

$$E = \frac{kQr}{R^3}$$

$$\frac{q_{in}}{Q} = \frac{\frac{4}{3}\pi r^3}{\frac{4}{3}\pi R^3}$$

$$q_{in} = Q \frac{r^3}{R^3}$$

3. A thin spherical shell of radius R has a total charge Q distributed uniformly over its surface. Find the E-field at


a) points outside the shell

$$E = \frac{kQ}{r^2}$$

b) inside the shell

$$E = 0, q_{in} = 0$$

4. Find the electric field a distance r from a line of positive charge of infinite length constant charge per unit length λ .



5. Find the E-field due to an insulating, infinite plane of positive charge with uniform surface charge density σ .

$$\oint E \cdot dA = \frac{q_{in}}{\epsilon_0}$$


$$q_{in} = \sigma A$$



$$E(2A) = \frac{\sigma A}{\epsilon_0}$$

$$\sigma = \frac{q}{A}$$

$$E = \frac{\sigma}{2\epsilon_0}$$

Sheet





4. $\oint E \cdot dA = \frac{q_{in}}{\epsilon_0}$
 $E (2\pi r l) = \frac{\lambda l}{\epsilon_0}$
 $E = \frac{\lambda}{2\pi \epsilon_0 r} = \frac{2k\lambda}{r}$